Name:	
(5/11/17)	



Summer Packet Pre-AP Algebra 1- 2018-19

To receive credit all work must be shown. There will be a test the first week back from summer on this packet. Any work done on additional paper must be turned in with the assignment.

A. Order of Operations

PEMDAS = Parentheses, Exponents, Multiplication/Division, Add/Subtract from left to right.

Simplify each expression using appropriate Order of Operations.

- 1. $1 \bullet 5 6 \div 2 + 3^2$ 3. $4 + 2(10 4 \bullet 6)$ 5. $12(20 17) 3 \bullet 6$
- 2. $125 \div [5(2+3)]$ 4. $3(2+7)^2 \div 5$ 6. $3^2 \div 3 + 2^2 \bullet 7 - 20 \div 5$

The fraction bar represents division: $\frac{1}{2} = 1 \div 2 = 0.5$ $1 \div 4 = \frac{1}{4} = 0.25$ Fractions should always be written in simplest form: $\frac{3}{30} = \frac{3 \cdot 1}{3 \cdot 10} = \frac{1}{10}$ $\frac{5}{20} = \frac{1 \cdot 5}{4 \cdot 5} = \frac{1}{4}$ Any integer can be written as fraction with a denominator of 1: $5 = \frac{5}{1}$ $-8 = -\frac{8}{1}$ $-32 = -\frac{32}{1}$ An improper fraction can be written as a mixed number (but improper fractions are more useful so don't convert): $\frac{17}{5} = \frac{15+2}{5} = \frac{15}{5} + \frac{2}{5} = 3 + \frac{2}{5} = 3\frac{2}{5} \qquad \qquad \frac{25}{3} = \frac{24+1}{3} = 8\frac{1}{3}$ There are several equivalent ways to write a negative fraction: $-\frac{3}{5} = \frac{-3}{5} = \frac{3}{-5} \qquad -\frac{7}{13} = \frac{-7}{13} = \frac{7}{-13} \qquad \frac{-3}{-5} = \frac{3}{5}$ To add or subtract fractions, you must have a common denominator: $\frac{1}{5} + \frac{2}{15} = \frac{3}{15} + \frac{2}{15} = \frac{5}{15} = \frac{1}{3}$ $\frac{3}{10} + \frac{1}{6} = \frac{9}{30} + \frac{5}{30} = \frac{14}{30} = \frac{7}{15}$ To multiply fractions, multiply the numerators and the denominators: $\frac{1}{3} \cdot \frac{2}{15} = \frac{2}{45} \qquad \qquad \frac{3}{7} \cdot \frac{4}{9} = \frac{12}{63} = \frac{4}{21}$ To divide fractions, multiply by the reciprocal: $\frac{2}{7} \div \frac{10}{21} = \frac{2}{7} \cdot \frac{21}{10} = \frac{1}{1} \cdot \frac{3}{5} = \frac{3}{5} \qquad \qquad \frac{6}{5} \div \frac{9}{8} = \frac{6}{5} \cdot \frac{8}{9} = \frac{48}{45} = \frac{16}{15}$

Evaluate the following expressions and record each answer as a fraction in simplest form:

1.
$$\frac{1}{6} + \frac{5}{18} =$$
 4. $\frac{3}{7} \div \frac{5}{8} =$

2.
$$\frac{1}{5} - \frac{2}{15} =$$
 5. $\frac{-4}{3} \div \frac{3}{5} =$

3.
$$\frac{7}{10} + \frac{5}{8}$$
 6. $\frac{14}{26} - \frac{6}{13} =$

7.
$$\frac{9}{2} \cdot \frac{-4}{3} =$$
 9. $4\frac{1}{5} \div \frac{3}{5} =$


8.
$$\frac{3}{-4} \cdot \frac{-2}{9} =$$
 10. $\left(4\frac{1}{5}\right) \left(\frac{3}{5}\right) =$

C. Coordinate Plane

Label : the origin, the 4 quadrants, the x-axis, and the y-axis

Plot and label the ordered pairs:

A: (3,-4)	B: (-2,1)	C: (6.5, 0)	D: (2,9 ¹ / ₂)
E: (0, 3)	F: (-4, - ⁹ / ₂)	<i>G</i> : (8,-2)	H: (- ¹⁶ / ₂ , 2)



		D.	Integer Operatio	ons	
1.	-23 + 17 =				-15
2.	15 - (-6) =			7.	$\frac{-15}{-3}$
3.	-5 - 8 =			8.	-24 + 30 + -9 =
4.	12 * -3 =				
5.	-4 * -7 =			9.	15 + 12 - 8 + 5 =
6.	$\frac{18}{-6}$				

E. Algebraic Properties

Write an example of each property:

- Commutative Property of Addition
 Commutative Property of Multiplication
 Associative Property of Addition
 Associative Property of Multiplication
 Additive Inverse
 Additive Identity
- 7. Multiplicative Inverse 8. Multiplicative Identity
- 9. Multiplicative Property of Zero 10. Distributive Property

Like terms: terms that have the same variables with the same exponents
CLT: add the coefficients of the like terms.

2x + 5x = 7x $3x^{2} - 7x + 8x^{2} + 8 = 11x^{2} - 7x + 8$ $6xy + 7x^{2} - 8xy - 9x + 10x^{2} = 17x^{2} - 9x - 2xy$

Simplify by combining like terms:

1.
$$6n + 11n$$
 2. $6x^2 - 7x + 9 - 8x^2 + 7x - 8$ 3. $8xy - 2x + 9xy + 3x$

- 4. $25x^2 9x + 7y 13x^2 + 8y$ 5. $8x^2y + 9xy - 2x^2y + 3xy - 9xy^2$
- 6. 6a⁴b 7ab + 3b 6a⁴b + 7ab

G. Distributive Property

Simplify using the Distributive Property:

1. -2(3y - 7) 2. 6(1 - 4x)

- 3. 4(2x + 5) 4. -3(5x 10 + 7y)
- 5. -(4x 5y) 6. 2x(-3x 1)

7. -x(5x + 2 - 7y)

H. Evaluating With Integers

EVALUATE (simplify) the expression using your order of operations and integer rules. Show the substitution step and all work. Reduce all fractions. Use the following given values to evaluate the following expressions.

a = -2	b = 2	c = -3	d = 4	e = -5	f = -1
1. 5a + b - 2c			5. $\frac{2(b-e)}{c+d}$		
$\frac{2e}{a} - ab + f$			62f(a + 20	c)	
$\frac{(3b-c)}{(6c+c)}$			73d + af -	- 15	
42a - 4d - f			8. (3f - c)(2	a - b)(-d)	


I.	Sal	wina	Ea	uations
±.	30	ving	CY	uu 110/13

	1. Solving Equation	15	
Find the numeric value	of the variable by isolating the varia	ble.	
🗸 Inverse Operati	ons cancel each other:		
addition \leftrightarrow su	btraction or multiplication \leftrightarrow division	or square \leftrightarrow square root	
✓ Apply the invers	e operation to both sides of the equation	on to isolate the variable	
✓ Always balance a	cross the equal sign.		
Solve for the variable in	each problem.		
1. x - 4 = 16	3. a - 4 = 15	5. 7x = 42	

2. 25 + x = 17

4. -8m = 64


6. 4x + 7 = 31




J. Solving Inequalities

<u>K.</u> Perimeter To find the perimeter (distance around) any shape, add all of the sides.

Find the perimeter of the following figures. Round to the nearest hundredth if necessary.






3 ft

L. Area

Triangle:	$A = rac{1}{2}bh$ where b is the length of the base and h is the height of the triangle.
Square:	$A = s^2$
Parallelogram/F	Rectangle: $A=bh$ where b is the length of the base and h is the height
Trapezoid:	$A=rac{1}{2}hig(b_1^{}+b_2^{}ig)$ where h is the height, and b1 and b2 are the bases
Circle:	$A=\pi r^2$ where r is the radius of the circle


Find the area of the following figures. Round to the nearest hundredth if necessary.



M. Surface Area


Prism:	$Surface \ Area = Ph + 2B; Volume = Bh \$ where P = Perimeter of base, h = height of prism B = Area of base
Cylinder:	Surface $Area=2\pi rh+2\pi r^2;$ $Volume=\pi r^2h$ where r = radius of cylinder h = height of cylinder
Pyramid:	Surface $Area = \frac{1}{2}P + B$; $Volume = \frac{1}{3}Bh$ where P = Perimeter of base, =slant height, h = height of pyramid
	B = Area of base
Cone:	Surface Area = $\pi r + \pi r^2$; Volume = $\frac{1}{3}\pi r^2 h$ where r = radius of cone, h = height of cone, =slant height

Find the surface area of the following figures. Round to the nearest hundredth if necessary.



_		
	Prism:	<i>V = Bh</i> where P = Perimeter of base, h = height of prism B = Area of base
	Cylinder:	$V = \pi r^2 h$ where r = radius of cylinder h = height of cylinder
	Pyramid: B = Area of	<i>V = (1/3)Bh</i> where P = Perimeter of base, =slant height, h = height of pyramid f base
	Cone:	$V = (1/3)\pi r^2 h$ where r = radius of cone, h = height of cone, =slant height

Find the volume of the following figures. Round to the nearest hundredth if necessary.







Slope - m, rate of change,
$$\frac{rise}{run}$$
 , $m = \frac{y_2 - y_1}{x_2 - x_1}$

Find the slope of the line that passes through the points.

1.




2.




(-3, 3	,	y	(2,	3)	+
	-1-				
-3	-1-1-	1	_	3	$\frac{1}{x}$
	,	,			

3.

N. X- and Y-Intercepts



Find the x- and y-intercepts for the graphs below.



O. Slope-Intercept Form

Slope-Intercept Form: y = mx + b, where m is the slope of the line, and b is the y-intercept.


Write and equation of a line with the given slope and y-intercept.

1. Slope: 7; y-intercept: 4

2. Slope: -3, y-intercept: 5


3. Slope: 1, y-intercept: -6

4.



P. Linear vs. Nonlinear Functions

You can determine if a function is linear or nonlinear by looking at the rate of change.



As x increases by 2, y increases by 3. The rate of change is constant, so this function is linear.

As x increases by 1, y decreases by a different amount each time. The rate of change is not constant, so this function is nonlinear.

Determine whether each table represents a linear or nonlinear function.

1.

х	3	5	7	9
У	7	9	11	13

X	1	5	9	13
у	0	6	8	9

3.

х	3	6	9	12
у	2	3	4	5


4	4	•

2.

Х	-2	-3	-4	-5
У	-1	-5	9	8


Q. Pythagorean Theorem

Using Pythagorean Theorem $a^2 + b^2 = c^2$ (where a and b are the legs, and c is the hypotenuse), find the missing lengths.



R. Functions

A function relates each element of a set with exactly one element of another set. Summary \rightarrow For every input, there is one and only one output.



Determine if the following are functions or not.

1. {(0, 0), (1, 1), (1, -1), (2, 2), (2, -2)}

2. {(-2, 2), (-1, 1), (0, 0), (1, 1), (2, 2)}

3.	x	4	2	5	7
	У	6	4	5	8

4.	x	1	2	3	2
	у	4	5	6	7