LISD Allergy Action Plan for Elementary Students Place D.O.B ____/___ Student's Campus: _____ Grade: ____ Teacher____ **Picture** Severe Allergy to: Here **Asthma:** Yes (higher risk for a severe reaction) Weight _____ lbs. Student history and warning signs: MILD SYMPTOMS **SEVERE SYMPTOMS** a few hives, mild itching many hives all over, redness, swelling of face, eyes, or lips Mouth: itchy mouth Lung: short of breath, wheezing, repetitive cough Stomach: mild nausea or discomfort Throat: tight, hoarse, trouble breathing or swallowing itchy, runny nose, sneezing Mouth: swelling of tongue and/or lips Stomach: vomiting, diarrhea, severe cramping TREATMENT PLAN Heart: pale, blue, faint, weak pulse, dizzy, confusion, loss of consciousness (TWO CHOICES – PLEASE CHECK ONLY ONE): Others: anxiety, feeling bad, or feeling of impending doom ORDERED MEDICATIONS AND DOSES Plan 1: For MILD SYMPTOMS: Antihistamine Brand: [] Benadryl or Diphenhydramine Mild symptoms from MORE THAN ONE BODY AREA (skin, mouth, stomach, or [] Other: _ nose) are TREATED AS <u>SEVERE</u> SYMPTOMS!!! Give EPINEPHRINE. Antihistamine Dose: Mild Symptoms from a **single** body area: [] 12.5 mg [] 18.75 mg [] 25 mg 1. Give Antihistamine if ordered. [] 31.25 mg [] 37.5 mg [] 43.75 mg 2. Stay with student and monitor for worsening symptoms. 3. If symptoms progress, USE EPINEPHRINE (treat as SEVERE symptoms). [] 50 mg OTHER ____mg 4. Contact parent. Nurses Notes: _____mg = ____ For SEVERE SYMPTOMS: 1. ADMINISTER EPHINEPHRINE IMMEDIATELY. **EPINEPHRINE Dose:** Call 911. INJECTION [] 0.15 mg IM [] 0.3 mg IM Give *Antihistamine* and then **Inhaler** if ordered (and not already used). INHALATION [] 1 mg [] 2 mg 4. Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side. [] If not improved, give second dose of Epinephrine in If symptoms do not improve, or return, more epinephrine may be needed. See order if you need to repeat the dose and when dose is to be repeated. ____ minutes. Contact parent. 1 Student will not have second dose of Epinephrine at school. _____ Parent's Initials OR Type of injector____ Plan 2: Give Epinephrine immediately for ANY symptoms if the allergen was likely eaten: Inhaler or Other (e.g., inhaler-bronchodilator if asthmatic): 1. ADMINISTER EPHINEPHRINE IMMEDIATELY. Call 911. 2. Give Antihistamine and then Inhaler if ordered. Dosage: _____ Route: _____ Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side. If symptoms do not improve, or return, more epinephrine may be needed. Frequency: See order if you need to repeat the dose and when dose is to be repeated. Indication for use: _____ Contact parent. I request and authorize Lewisville ISD personnel to administer the above medication as prescribed. I understand that the school administrator may designate any qualified person or persons to administer these medications. This form is valid for one school year. Physician must be licensed to practice in Texas. Temporary (2 months) orders for out of state US Physicians are acceptable to initiate treatment for transferring students. A signature is required to authorize the registered nurse and the prescribing physician to discuss and/or clarify the medication order and the student's response to the treatment plan. Elementary students are not permitted to transport medications. Unused medications not picked up at the end of the school year will be disposed of properly. Printed Name: Physician Signature: Parent Signature: _____ Date: _____ Office #: _____ ____ Fax #: ____ Address:

Rev. 5/2025 Epinephrine Expires: _____ Lot #: ____ Antihistamine Expires: ____ Inhaler Expires: ____ Inhaler Expires: ____