Name: Grade: Grade:						Place Student's Picture
evere A	llergy to:					Here
sthma:	∐Yes (h	higher risk for a severe reaction)	🗌 No	w	eight lbs.	
		MILD SYMPTOMS		SEVERE SYMPTOMS		
ЕАТМЕ УО СНОІ	Skin: Mouth: Stomach: Nose: NT PLAN CES – PLE	a few hives, mild itching itchy mouth mild nausea or discomfort itchy, runny nose, sneezing ASE CHECK ONLY <u>ONE</u> :	Skin: Lung: Throat: Mouth: Stomach: Heart: Others:	short of breat tight, hoarse, swelling of tor vomiting, diar pale, blue, fai	Il over, redness, swelling of face, eyes, or lips h, wheezing, repetitive cough trouble breathing or swallowing ngue and/or lips rhea, severe cramping nt, weak pulse, dizzy, confusion, loss of conse ig bad, or feeling of impending doom	
Plai		For <u>MILD SYMPTOMS</u> :			ORDERED MEDICATIONS Antihistamine Brand: [] Benadryl or Diphenhydrami	
 nose) are TREATED AS <u>SEVERE</u> SYMPTOMS!!! Give EPINEPHRINE. Mild Symptoms from a single body area: Give Antihistamine if ordered. Stay with student and monitor for worsening symptoms. If symptoms progress, USE EPINEPHRINE (treat as SEVERE symptoms). Contact parent. For <u>SEVERE SYMPTOMS</u>: 			Antihistamine Dose: [] 12.5 mg [] 18.75 mg [] 31.25 mg [] 37.5 mg [] 50 mg Nurses Notes:mg =	[] 43.75 mg		
 INJECT EPHINEPHRINE IMMEDIATELY. Call 911. Give Antihistamine and then Inhaler if ordered (and not already used). Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side. If symptoms do not improve, or return, more epinephrine may be needed. See order if you need to repeat the dose and when dose is to be repeated. Contact parent. 					EPINEPHRINE Dose: [] 0.15 mg IM [] 0.15 EPINEPHRINE Brand: [] 1 [] EpiPen [] 1 [] If not improved, give second	Auvi-Q
OR					[] Student will not have second	dose of Epinephrine
aller 1. 2. (3. (4. 5. 5.	rgen was li INJECT EPI Call 911. Give <i>Antihi</i> s Lay the pers are vomiting If symptoms	E Epinephrine immediately for a kely eaten: HINEPHRINE IMMEDIATELY. stamine and then Inhaler if ordered. son flat, raise legs and keep warm. If l, let them sit up or lie on their side. do not improve, or return, more epin you need to repeat the dose and whe ent.	breathing is difficul	It or they eded.	at school. Pa Inhaler or Other (e.g., inhaler-bronchodilator Brand: Dosage: Dosage: Pa Frequency: Pa Indication for use: Pa	if asthmatic):

Elementary students are not permitted to transport medications. Onused medications not picked up at the end of the school year will be disposed of property.								
Physician Signature:	Printed Name:	Parent Signature:						
Date: 0	Office #: Fax #:	Date:						
Address:								

Revised 4/17