LISD Allergy Action Plan for Secondary Students

Name: D.O.B/				Place Student's	
Campus:	Grade:	Teacher		Picture	
Severe Allergy to:				Here	
Student history and warning signs:				-	
Does Student Have Asthma?Yes _	No				

N	MILD SYMPTOMS				
Skin:	a few hives, mild itching				
Mouth:	itchy mouth				
Stomach:	mild nausea or discomfort				
Nose:	itchy, runny nose, sneezing				

SEVERE SYMPTOMS Skin: many hives all over, redness, swelling of face, eyes, or lips Lung: short of breath, wheezing, repetitive cough tight, hoarse, trouble breathing or swallowing Throat: Mouth: swelling of tongue and/or lips Stomach: vomiting, diarrhea, severe cramping Heart: pale, blue, faint, weak pulse, dizzy, confusion, loss of consciousness Others: anxiety, feeling bad, or feeling of impending doom

ORDERED MEDICATIONS AND DOSES

Student may self-carry & administer medications and has

TREATMENT PLAN (TWO CHOICES - PLEASE CHECK ONLY ONE):

Plan 1:	For MILD SYMPTOMS:

Mild symptoms from MORE THAN ONE BODY AREA (skin, mouth, stomach, or nose) are TREATED AS <u>SEVERE</u> SYMPTOMS!!! Give EPINEPHRINE.

Mild Symptoms from a single body area:

- 1. Give Antihistamine if ordered.
- 2. Stay with student and monitor for worsening symptoms.
- 3. If symptoms progress, USE EPINEPHRINE (treat as SEVERE symptoms).
- 4. Contact parent.

For <u>SEVERE SYMPTOMS</u>:

- 1. INJECT EPHINEPHRINE IMMEDIATELY.
- Call 911.
- Give Antihistamine and then Inhaler if ordered (and not already used).
- 4. Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
- If symptoms do not improve, or return, more epinephrine may be needed. See order if you need to repeat the dose and when dose is to be repeated.
- 6. Contact parent.

OR

Plan 2: Give Epinephrine immediately for ANY symptoms if the allergen was likely eaten:

- INJECT EPHINEPHRINE IMMEDIATELY. 1.
- Call 911.
- 3. Give Antihistamine and then Inhaler if ordered.
- 4. Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
- If symptoms do not improve, or return, more epinephrine may be needed. See order if you need to repeat the dose and when dose is to be repeated.
- 6. Contact parent.

been instructed on proper use Y N
Antihistamine Brand: [] Benadryl or Diphenhydramine [] Other:
Antihistamine Dose: [] 12.5 mg
[] 31.25 mg
[] 50 mg Nurses Notes:mg =
Medication is kept:with studentin clinic both
EPINEPHRINE Dose: [] 0.15 mg IM [] 0.3 mg IM
Type of injector:
[] If not improved, give second dose of Epinephrine in minutes. [] Student will not have second dose of Epinephrine at school Parent's Initials Medication is kept:with student in clinic both
Inhaler:
Brand:
Dosage: Route:
Frequency:
Indication for use:with studentin clinicboth

I request and authorize Lewisville ISD personnel to administer the above medication as prescribed. I understand that the school administrator may designate any qualified person or persons to administer these medications. This form is valid for one school year. Physician must be licensed to practice in Texas. Temporary (2 months) orders for out of state US Physicians are acceptable to initiate treatment for transferring students. A signature is required to authorize the registered nurse and the prescribing physician to discuss and/or clarify the medication order and the student's response to the treatment plan. Elementary students are not permitted to transport medications. Unused medications not picked up at the end of the school year will be disposed of properly.

	Physician Signature:		Printed Name:	Parent Signature:
	Date:	Office #:	Fax #:	Date:
	Address:			
_	euland 4/47 Eninophrin	o Evniroci	Let # Panadryl Expires	Inhalar Evniros: