

Name: \_\_\_\_\_

# Hebron Ninth Grade Campus Algebra 1 Honors Summer Packet

We have selected some middle school math topics that will help you be better prepared for Algebra 1 Honors. Try not to use calculator. Show all work. See how much you remember. This packet is optional and is not for a grade.

## A. Order of Operations

PEMDAS = Parentheses, Exponents, Multiplication/Division, Add/Subtract from left to right.

Simplify each expression using appropriate Order of Operations.

1.  $1 \cdot 5 - 6 \div 2 + 3^2$

3.  $4 + 2(10 - 4 \cdot 6)$

5.  $12(20 - 17) - 3 \cdot 6$

2.  $125 \div [5(2 + 3)]$

4.  $3(2 + 7)^2 \div 5$

6.  $3^2 \div 3 + 2^2 \cdot 7 - 20 \div 5$

## B. Fractions Review

The fraction bar represents division:

$$\frac{1}{2} = 1 \div 2 = 0.5 \qquad 1 \div 4 = \frac{1}{4} = 0.25$$

Fractions should always be written in simplest form:

$$\frac{5}{20} = \frac{1 \cdot 5}{4 \cdot 5} = \frac{1}{4} \qquad \frac{3}{30} = \frac{3 \cdot 1}{3 \cdot 10} = \frac{1}{10}$$

Any integer can be written as fraction with a denominator of 1:

$$5 = \frac{5}{1} \qquad -8 = -\frac{8}{1} \qquad -32 = -\frac{32}{1}$$

An improper fraction can be written as a mixed number (but improper fractions are more useful so don't convert):

$$\frac{17}{5} = \frac{15+2}{5} = \frac{15}{5} + \frac{2}{5} = 3 + \frac{2}{5} = 3\frac{2}{5} \qquad \frac{25}{3} = \frac{24+1}{3} = 8\frac{1}{3}$$

There are several equivalent ways to write a negative fraction:

$$-\frac{3}{5} = \frac{-3}{5} = \frac{3}{-5} \qquad -\frac{7}{13} = \frac{-7}{13} = \frac{7}{-13} \qquad \frac{-3}{-5} = \frac{3}{5}$$

To add or subtract fractions, you must have a common denominator:

$$\frac{1}{5} + \frac{2}{15} = \frac{3}{15} + \frac{2}{15} = \frac{5}{15} = \frac{1}{3} \qquad \frac{3}{10} + \frac{1}{6} = \frac{9}{30} + \frac{5}{30} = \frac{14}{30} = \frac{7}{15}$$

To multiply fractions, multiply the numerators and the denominators:

$$\frac{1}{3} \cdot \frac{2}{15} = \frac{2}{45} \qquad \frac{3}{7} \cdot \frac{4}{9} = \frac{12}{63} = \frac{4}{21}$$

To divide fractions, multiply by the reciprocal:

$$\frac{2}{7} \div \frac{10}{21} = \frac{2}{7} \cdot \frac{21}{10} = \frac{1}{1} \cdot \frac{3}{5} = \frac{3}{5} \qquad \frac{6}{5} \div \frac{9}{8} = \frac{6}{5} \cdot \frac{8}{9} = \frac{48}{45} = \frac{16}{15}$$

Evaluate the following expressions and record each answer as a fraction in simplest form:

1.  $\frac{1}{6} + \frac{5}{18} =$

4.  $\frac{3}{7} \div \frac{5}{8} =$

2.  $\frac{1}{5} - \frac{2}{15} =$

5.  $\frac{-4}{3} \div \frac{3}{5} =$

3.  $\frac{7}{10} + \frac{5}{8} =$

6.  $\frac{14}{26} - \frac{6}{13} =$

$$7. \frac{9}{2} \cdot \frac{-4}{3} =$$

$$9. 4\frac{1}{5} \div \frac{3}{5} =$$

$$8. \frac{3}{-4} \cdot \frac{-2}{9} =$$

$$10. \left(4\frac{1}{5}\right)\left(\frac{3}{5}\right) =$$

### C. Coordinate Plane

Label : the origin, the 4 quadrants, the x-axis, and the y-axis

Plot and label the ordered pairs:

A: (3, -4)

B: (-2, 1)

C: (6.5, 0)


D:  $(2, 9\frac{1}{2})$

E: (0, 3)

F:  $(-4, -\frac{9}{2})$

G: (8, -2)

H:  $(-\frac{16}{2}, 2)$



### D. Integer Operations

1.  $-23 + 17 =$

2.  $15 - (-6) =$

3.  $-5 - 8 =$

4.  $12 * -3 =$

5.  $-4 * -7 =$

6.  $\frac{18}{-6}$

7.  $\frac{-15}{-3}$

8.  $-24 + 30 + -9 =$

9.  $15 + 12 - 8 + 5 =$

### E. Combining Like Terms

- Like terms: terms that have the same variables with the same exponents
- CLT: add the coefficients of the like terms.

$$2x + 5x = 7x \quad 3x^2 - 7x + 8x^2 + 8 = 11x^2 - 7x + 8 \quad 6xy + 7x^2 - 8xy - 9x + 10x^2 = 17x^2 - 9x - 2xy$$

Simplify by combining like terms:

1.  $6n + 11n$

2.  $6x^2 - 7x + 9 - 8x^2 + 7x - 8$

3.  $8xy - 2x + 9xy + 3x$

4.  $25x^2 - 9x + 7y - 13x^2 + 8y$

5.  $8x^2y + 9xy - 2x^2y + 3xy - 9xy^2$

6.  $6a^4b - 7ab + 3b - 6a^4b + 7ab$

## F. Distributive Property

Simplify using the Distributive Property:

1.  $-2(3y - 7)$

2.  $6(1 - 4x)$

3.  $4(2x + 5)$

4.  $-3(5x - 10 + 7y)$

5.  $-(4x - 5y)$

6.  $2x(-3x - 1)$

7.  $-x(5x + 2 - 7y)$

## G. Evaluating With Integers

EVALUATE (simplify) the expression using your order of operations and integer rules. Show the substitution step and all work. Reduce all fractions. Use the following given values to evaluate the following expressions.

$a = -2$

$b = 2$

$c = -3$

$d = 4$

$e = -5$

$f = -1$

1.  $5a + b - 2c$

5.  $\frac{2(b - e)}{c + d}$

2.  $\frac{2e}{a} - ab + f$

6.  $-2f(a + 2c)$

3.  $\frac{(3b - c)}{(6c + c)}$

7.  $-3d + af - 15$

$$4. -2a - 4d - f$$

$$8. (3f - c)(2a - b)(-d)$$

### H. Solving Equations

Find the numeric value of the variable by isolating the variable.

- ✓ Inverse Operations cancel each other:  
addition  $\leftrightarrow$  subtraction or multiplication  $\leftrightarrow$  division or square  $\leftrightarrow$  square root
- ✓ Apply the inverse operation to both sides of the equation to isolate the variable
- ✓ Always balance across the equal sign.

Solve for the variable in each problem.

$$1. x - 4 = 16$$

$$3. a - 4 = 15$$

$$5. 7x = 42$$

$$2. 25 + x = 17$$

$$4. -8m = 64$$

$$6. 4x + 7 = 31$$

## I. Solving Inequalities


Symbol	Meaning	Equation or Inequality	Graph
=	equals	$x = 3$	
<	is less than	$x < 3$	
≤	is less than or equal to	$x ≤ 3$	
>	is greater than	$x > 3$	
≥	is greater than or equal to	$x ≥ 3$	

**Examples:**

$$2x \leq 4$$

$$x \leq 2$$

Divide each side by 2



$$-4y < 18$$

$$\frac{-4y}{-4} > \frac{18}{-4}$$

$$y > -4.5$$


Divide by -4 and change < to >

Simplify




Solve and graph the following inequalities.

1.  $3f < 15$



2.  $m + 6 \geq 7$



3.  $-7h < 56$



4.  $2g - 8 > 20$







5.  $3x + 5 < -22$



### J. Perimeter

To find the perimeter (distance around) any shape, add all of the sides.

Find the perimeter of the following figures. Round to the nearest hundredth if necessary.



### K. Area

**Triangle:**  $A = \frac{1}{2}bh$  where  $b$  is the length of the base and  $h$  is the height of the triangle.



**Square:**  $A = s^2$

**Parallelogram/Rectangle:**  $A = bh$  where  $b$  is the length of the base and  $h$  is the height


**Trapezoid:**  $A = \frac{1}{2}h(b_1 + b_2)$  where  $h$  is the height, and  $b_1$  and  $b_2$  are the bases

**Circle:**  $A = \pi r^2$  where  $r$  is the radius of the circle


Find the area of the following figures. Round to the nearest hundredth if necessary.




3.







5.



### L. Surface Area

**Prism:**  $Surface\ Area = Ph + 2B$ ;  $Volume = Bh$  where  $P$  = Perimeter of base,  $h$  = height of prism  $B$  = Area of base





**Cylinder:**  $Surface\ Area = 2\pi rh + 2\pi r^2$ ;  $Volume = \pi r^2 h$  where  $r$  = radius of cylinder  $h$  = height of cylinder

**Pyramid:**  $Surface\ Area = \frac{1}{2}P\ell + B$ ;  $Volume = \frac{1}{3}Bh$  where  $P$  = Perimeter of base,  $\ell$  = slant height,  $h$  = height of pyramid  
 $B$  = Area of base

**Cone:**  $Surface\ Area = \pi r\ell + \pi r^2$ ;  $Volume = \frac{1}{3}\pi r^2 h$  where  $r$  = radius of cone,  $h$  = height of cone,  $\ell$  = slant height

Find the **surface area** of the following figures. Round to the nearest hundredth if necessary.

1.



2.

3.

4.

### M. Volume

**Prism:**  $V = Bh$  where  $P$  = Perimeter of base,  $h$  = height of prism  $B$  = Area of base


**Cylinder:**  $V = \pi r^2 h$  where  $r$  = radius of cylinder  $h$  = height of cylinder

**Pyramid:**  $V = (1/3)Bh$  where  $P$  = Perimeter of base,  $\ell$  = slant height,  $h$  = height of pyramid  
 $B$  = Area of base


**Cone:**  $V = (1/3)\pi r^2 h$  where  $r$  = radius of cone,  $h$  = height of cone,  $\ell$  = slant height

Find the **volume** of the following figures. Round to the nearest hundredth if necessary.


5.




6.



7.



8.




### N. Slope


**Slope -  $m$ ,** rate of change,  $\frac{\text{rise}}{\text{run}}$ ,  $m = \frac{y_2 - y_1}{x_2 - x_1}$

Find the slope of the line that passes through the points.


1.



2.




3.



### O. X- and Y-Intercepts

**X-Intercept:** the point where the line intersects the x (horizontal)-axis. The y value of the point will be 0.




**Y-Intercept:** the point where the line intersects the y (vertical)-axis. The x value of the point will be 0.



For the graph pictured to the left, the x-intercept would be located at A(-2, 0) and the y-intercept would be located at B(2, 0).

Find the x- and y-intercepts for the graphs below.

1.



3.

2.

### P. Slope-Intercept Form

**Slope-Intercept Form:**  $y = mx + b$ , where  $m$  is the slope of the line, and  $b$  is the y-intercept.


Write an equation of a line with the given slope and y-intercept.

1. Slope: 7; y-intercept: 4



2. Slope: -3; y-intercept: 5

3. Slope: 1; y-intercept: -6

4.



5.



6.

### Q. Linear vs. Nonlinear Functions

You can determine if a function is linear or nonlinear by looking at the rate of change.

		+2	+2	+2
		↘	↘	↘
<b>x</b>	3	5	7	9
<b>y</b>	7	10	13	16
		↗	↗	↗
		+3	+3	+3

As  $x$  increases by 2,  $y$  increases by 3. The rate of change is constant, so this function is linear.

		+1	+1	+1
		↘	↘	↘
<b>x</b>	1	2	3	4
<b>y</b>	-3	-6	-10	-15
		↗	↗	↗
		-3	-4	-5

As  $x$  increases by 1,  $y$  decreases by a different amount each time. The rate of change is not constant, so this function is nonlinear.

Determine whether each table represents a linear or nonlinear function.

1.

<b>x</b>	3	5	7	9
<b>y</b>	7	9	11	13

2.

<b>x</b>	1	5	9	13
<b>y</b>	0	6	8	9

3.

<b>x</b>	3	6	9	12
<b>y</b>	2	3	4	5


4.

<b>x</b>	-2	-3	-4	-5
<b>y</b>	-1	-5	9	8


### R. Pythagorean Theorem

Using Pythagorean Theorem  $a^2 + b^2 = c^2$  (where  $a$  and  $b$  are the legs, and  $c$  is the hypotenuse), find the missing lengths.


1.




2.



3.




4.



### S. Functions

A function relates each element of a set with exactly one element of another set. Summary → For every input, there is one and only one output.



Determine if the following are functions or not.

1.  $\{(0, 0), (1, 1), (1, -1), (2, 2), (2, -2)\}$

2.  $\{(-2, 2), (-1, 1), (0, 0), (1, 1), (2, 2)\}$

3.

$x$	4	2	5	7
$y$	6	4	5	8

4.

$x$	1	2	3	2
$y$	4	5	6	7